

INSTRUCTIONS FOR YOUR LENS PROCEDURE

In preparation for your upcoming surgery, please review the following reminders:

General Reminders:

- A member of our team will call you the day before your procedure to review your medical history and to let know your surgery appointment time.
- o Please fill your eyedrop prescriptions if you have not already done so.

• Start using the following eyedrops <u>3 days</u> prior to your procedure:

- Place 1 DROP of <u>ILEVRO</u> 1 TIME EACH DAY and 1 DROP of <u>MOXIFLOXACIN (VIGAMOX)</u> 3 TIMES EACH DAY (breakfast, lunch, and bedtime) in your operative eye(s).
- Please wait <u>5 minutes</u> between eye drops.
- o **DO NOT USE DUREZOL**. You will start using this drop **after** your procedure.

• On the Morning of Your Procedure:

- o Place 1 DROP of <u>ILEVRO</u> and <u>MOXIFLOXACIN (VIGAMOX)</u> in your operative eye(s).
- Please wait <u>5 minutes</u> between eye drops.
- DO NOT USE DUREZOL. You will start using this drop after your procedure.
- You may take your regular medications in the morning with a sip of water (<u>AVOID</u> diabetic medications or diuretics).
- DO NOT apply cologne or perfume.
- DO NOT apply makeup or lotions to your face.
- Wear loose comfortable clothing: preferably a shirt or blouse that opens in the front. Leave jewelry and other valuables at home.
- HAVE A RESPONSIBLE ADULT ACCOMPANY AND DRIVE YOU TO THE CENTER AND BACK HOME.
- Notify the office of any change in your health (fever, cold, sore throat, etc.) before the procedure.

For patients having IV sedation:

- DO NOT eat or drink ANYTHING at least <u>8 HOURS</u> prior to your surgery. No gum or mints. You
 may brush your teeth, but do not swallow water.
- Please refrain from taking injectable GLP-1 agonists (Ozempic, Wegovy, Rybelsus, Mounjaro, Trulicity, Byetta, Victoza, Adlyxin, Bydureon, etc.) 1 week prior to your surgery date.

If you have any additional questions, please call SightTrust at (954) 653-0100.